


Claims

1. A compound of formula (1):



(1)

wherein:

----- is a single or double bond;

A is phenylene or heteroarylene;

m is 0, 1 or 2;

10 n is 0, 1 or 2;

R^1 is independently selected from halo, nitro, cyano, hydroxy, carboxy, carbamoyl, *N*- C_{1-4} alkylcarbamoyl, *N,N*-(C_{1-4} alkyl) $_2$ carbamoyl, sulphamoyl, *N*- C_{1-4} alkylsulphamoyl, *N,N*-(C_{1-4} alkyl) $_2$ sulphamoyl, -S(O) $_b$ C_{1-4} alkyl (wherein b is 0,1, or 2), C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, C_{1-4} alkanoyloxy, hydroxy C_{1-4} alkyl, fluoromethyl,

15 difluoromethyl, trifluoromethyl and trifluoromethoxy;

or, when n is 2, the two R^1 groups, together with the carbon atoms of A to which they are attached, may form a 4 to 7 membered ring, optionally containing 1 or 2 heteroatoms independently selected from O, S and N, and optionally being substituted by one or two methyl groups;


20 R^4 is independently selected from hydrogen, halo, nitro, cyano, hydroxy, fluoromethyl, difluoromethyl, trifluoromethyl, trifluoromethoxy, carboxy, carbamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy and C_{1-4} alkanoyl;

R^2 is hydrogen, hydroxy or carboxy;


R^3 is selected from hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkanoyl, carbamoyl, C_{3-7} cycloalkyl

25 (optionally substituted with 1 or 2 hydroxy groups), cyano(C_{1-4})alkyl, aryl, heterocyclyl, C_{1-4} alkyl (optionally substituted by 1 or 2 R^8 groups), and groups of the formulae B and B':

-78-



(B)



(B')

wherein y is 0 or 1, t is 0, 1, 2 or 3 and u is 1 or 2; provided that the hydroxy group is not a substituent on the ring carbon adjacent to the ring oxygen;

- 5 R^8 is independently selected from hydroxy, C_{1-4} alkoxy, C_{1-4} alkoxy, hydroxy C_{1-4} alkoxy, 5- and 6-membered cyclic acetals and mono- and di-methyl derivatives thereof, aryl, heterocyclyl, C_{3-7} cycloalkyl, C_{1-4} alkanoyl, C_{1-4} alkoxy, C_{1-4} alkylS(O)_b- (wherein b is 0, 1 or 2), C_{3-6} cycloalkylS(O)_b- (wherein b is 0, 1 or 2), arylS(O)_b- (wherein b is 0, 1 or 2), heterocyclylS(O)_b- (wherein b is 0, 1 or 2), benzylS(O)_b- (wherein b is 0, 1 or 2),
- 10 -N(OH)CHO, -C(=N-OH)NH₂, -C(=N-OH)NHC₁₋₄alkyl, -C(=N-OH)N(C₁₋₄alkyl)₂, -C(=N-OH)NHC₃₋₆cycloalkyl, -C(=N-OH)N(C₃₋₆cycloalkyl)₂, -COCOOR⁹, -C(O)N(R⁹)(R¹⁰), -NHC(O)R⁹, -C(O)NHSO₂(C₁₋₄alkyl), -NHSO₂R⁹, (R⁹)(R¹⁰)NSO₂-, -COCH₂OR¹¹, (R⁹)(R¹⁰)N- and -COOR⁹;
- R^9 and R^{10} are independently selected from hydrogen, hydroxy, C_{1-4} alkyl (optionally substituted by 1 or 2 R^{13}), C_{3-7} cycloalkyl (optionally substituted by 1 or 2 hydroxy groups), cyano(C_{1-4})alkyl, trihalo(C_{1-4})alkyl, aryl, heterocyclyl and heterocyclyl(C_{1-4} alkyl); or R^9 and R^{10} together with the nitrogen to which they are attached form a 4- to 6-membered ring where the ring is optionally substituted on carbon by 1 or 2 substituents independently selected from oxo, hydroxy, carboxy, halo, nitro, cyano, carbonyl, C_{1-4} alkoxy and
- 20 heterocyclyl; or the ring may be optionally substituted on two adjacent carbons by -O-CH₂-O- to form a cyclic acetal wherein one or both of the hydrogens of the -O-CH₂-O- group may be replaced by a methyl;
- R^{13} is selected from hydroxy, halo, trihalomethyl and C_{1-4} alkoxy;
- R^{11} is independently selected from hydrogen, C_{1-4} alkyl and hydroxy C_{1-4} alkyl;
- 25 or a pharmaceutically acceptable salt or pro-drug thereof.

2. A compound of the formula (1) as claimed in claim 1, wherein

R^3 is selected from hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkanoyl, carbamoyl, C_{3-7} cycloalkyl (optionally substituted with 1 or 2 hydroxy groups, cyano(C_{1-4})alkyl, phenyl, morpholino,

-79-

morpholinyl, piperidino, piperidyl, pyridyl, pyranyl, pyrrolyl, imidazolyl, thiazolyl, thienyl, thiadiazolyl, piperazinyl, isothiazolidinyl, 1,3,4-triazolyl, tetrazolyl, pyrrolidinyl, thiomorpholino, pyrrolinyl, homopiperazinyl, 3,5-dioxapiperidinyl, pyrimidyl, pyrazinyl, pyridazinyl, pyrazolyl, pyrazolinyl, isoxazolyl, 4-oxopyridyl, 2-oxopyrrolidinyl, 4-oxothiazolidyl, furyl, thienyl, oxazolyl, 1,3,4-oxadiazolyl, and 1,2,4-oxadiazolyl, tetrahydrothiopyranyl, 1-oxotetrahydrothiopyranyl, 1,1-dioxotetrahydrothiopyranyl and C₁₋₄alkyl (optionally substituted by 1 or 2 R⁸ groups);

R⁹ and R¹⁰ are independently selected from hydrogen, hydroxy, C₁₋₄alkyl (optionally substituted by 1 or 2 R¹³ groups), C₃₋₇cycloalkyl (optionally substituted by 1 or 2 hydroxy groups), cyano(C₁₋₄)alkyl, trihalo C₁₋₄alkyl, aryl, heterocyclyl and heterocyclyl(C₁₋₄alkyl);

or

R⁹ and R¹⁰ together with the nitrogen to which they are attached form a 4- to 6-membered ring where the ring is optionally substituted on carbon by 1 or 2 substituents selected from oxo, hydroxy, carboxy, halo, nitro, cyano, carbonyl and C₁₋₄alkoxy, or the ring may be optionally substituted on two adjacent carbons by -O-CH₂-O- to form a cyclic acetal wherein one or both of the hydrogens of the -O-CH₂-O- group may be replaced by a methyl;

R⁸ is independently selected from hydroxy, C₁₋₄alkoxyC₁₋₄alkoxy, hydroxyC₁₋₄alkoxy, 5- and 6-membered cyclic acetals and mono- and di-methyl derivatives thereof, aryl, heterocyclyl, C₃₋₇cycloalkyl, C₁₋₄alkanoyl, C₁₋₄alkoxy, C₁₋₄alkylS(O)_b- (wherein b is 0, 1 or 2), C₃₋₆cycloalkylS(O)_b- (wherein b is 0, 1 or 2), arylS(O)_b- (wherein b is 0, 1 or 2), heterocyclylS(O)_b- (wherein b is 0, 1 or 2), benzylS(O)_b- (wherein b is 0, 1 or 2), -N(OH)CHO, -C(=N-OH)NH₂, -C(=N-OH)NHC₁₋₄alkyl, -C(=N-OH)N(C₁₋₄alkyl)₂, -C(=N-OH)NHC₃₋₆cycloalkyl, -C(=N-OH)N(C₃₋₆cycloalkyl)₂, -COCOOR⁹, -C(O)N(R⁹)(R¹⁰), -NHC(O)R⁹, -C(O)NHSO₂(C₁₋₄alkyl), -NHSO₂R⁹, (R⁹)(R¹⁰)NSO₂-, -COCH₂OR¹¹, (R⁹)(R¹⁰)N- and -COOR⁹;

R¹³ is selected from hydroxy, halo, trifluoromethyl and C₁₋₄alkoxy;

R¹¹ is selected from hydrogen, C₁₋₄alkyl and hydroxyC₁₋₄alkyl;

or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

3. A compound of the formula (1) as claimed in claim 1 or claim 2 wherein:

R³ is selected from cyanoC₁₋₄alkyl and C₁₋₄alkyl (optionally substituted by 1 or 2 of R⁸ groups);

-80-

R^8 is independently selected from hydroxy, phenyl, 2,2-dimethyl-1,3-dioxolan-4-yl; 2,2-dimethyl-1,3-dioxan-4-yl; 2,2-dimethyl-1,3-dioxan-5-yl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, triazolyl, tetrazolyl, imidazolyl, pyrrolidinyl, piperidyl, tetrahydrofuryl, tetrahydropyranyl, tetrahydrothiopyranyl and tetrahydrothienyl, C_{1-4} alkoxy, C_{1-4} alkanoyl, $C_{1-4}alkylS(O)_b$ - (wherein b is 0, 1 or 2), $-C(O)N(R^9)(R^{10})$, $-COOR^9$, $-C(O)NHSO_2Me$, $-C(=N-OH)NH_2$, $-C(=N-OH)NHC_{1-4}alkyl$, $-C(=N-OH)N(C_{1-4}alkyl)_2$ and $-NHSO_2R^9$;

R^9 and R^{10} are independently selected from hydrogen, hydroxy, $C_{1-4}alkyl$ optionally substituted with R^{13} (wherein R^{13} is $C_{1-4}alkoxy$ or hydroxy); or

R^9 and R^{10} together with the nitrogen to which they are attached form a 4- to 6-membered ring where the ring may be optionally substituted on carbon by 1 or 2 hydroxy groups or carboxy groups), or the ring may be optionally substituted on two adjacent carbons by $-O-CH_2-O-$ to form a cyclic acetal wherein one or both of the hydrogens of the $-O-CH_2-O-$ group may be replaced by a methyl; or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

15

4. A compound of the formula (1) as claimed in any preceding claim, wherein:

R^3 is selected from cyano $C_{1-4}alkyl$ and $C_{1-4}alkyl$ (optionally substituted by 1 or 2 R^8 groups);

R^8 is independently selected from hydroxy, 2,2-dimethyl-1,3-dioxolan-4-yl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, tetrazolyl, $C_{1-4}alkoxy$, $C_{1-4}alkanoyl$, $C_{1-4}alkylS(O)_b$ - (wherein b is 0, 1 or 2), $-C(O)N(R^9)(R^{10})$, $-COOR^9$, $-C(O)NHSO_2Me$, $-C(=N-OH)NH_2$;

R^9 and R^{10} are independently selected from hydrogen, hydroxy, $C_{1-4}alkyl$ optionally substituted with R^{13} (wherein R^{13} is $C_{1-4}alkoxy$ or hydroxy); or

R^9 and R^{10} together with the nitrogen to which they are attached form a 4- to 6-membered ring selected from piperidine, 4-hydroxy piperidine, pyrrolidine, 3,4-dihydropyrrolidine and the dimethylacetal of 3,4-dihydropyrrolidine; or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

5. A compound of the formula (1) as claimed in any preceding claim, wherein: m is 1 and R^4 is chlorine; or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

6. A compound of the formula (1) as claimed in any preceding claim, wherein:

-81-

A is phenylene;

or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

7. A compound of the formula (1) as claimed in any one of claims 1 to 5, wherein:

5 A is heteroarylene;

or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

8. A compound of the formula (1) as claimed in any preceding claim, wherein:

---- is a single bond;

10 or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof.

9. A compound of the formula (1) as claimed in claim 1, which is any one of:

5-chloro-*N*-[1-(methoxycarbonylmethyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;

15 *N*-[1-(carboxymethyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-5-chloroindole-2-carboxamide;

5-chloro-*N*-(2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;

5-chloro-*N*-[2-oxo-1-[2-oxo-2-(pyridin-2-ylamino)ethyl]-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;

5-chloro-*N*-{1-[2-(methylthio)ethyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl}-1*H*-indole-2-

20 carboxamide;

5-chloro-*N*-{1-[2-(methylsulphinyl)ethyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl}-1*H*-indole-2-carboxamide;

5-chloro-*N*-{1-[2-(methylsulphonyl)ethyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl}-1*H*-indole-2-carboxamide;

25 5-chloro-*N*-{2-oxo-1-[2-oxo-2-(1,3,4-thiadiazol-2-ylamino)ethyl]-1,2,3,4-tetrahydroquinolin-3-yl}-1*H*-indole-2-carboxamide;

5-chloro-*N*-(1-{2-[(6-methylpyridin-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;

5-chloro-*N*-{2-oxo-1-[2-oxo-2-(pyridin-3-ylamino)ethyl]-1,2,3,4-tetrahydroquinolin-3-yl}-

30 1*H*-indole-2-carboxamide;

-82-

- 5-chloro-*N*-(1-{2-[(5-methyl-1,3,4-thiadiazol-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(5-ethyl-1,3,4-thiadiazol-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5 5-chloro-*N*-(1-{2-[(4-cyano-1*H*-pyrazol-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(4-methyl-1,3-thiazol-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(6-chloropyridin-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 10 5-chloro-*N*-(1-{2-[(3-hydroxypyridin-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(2-oxo-1-{2-oxo-2-[(pyridin-2-ylmethyl)amino]ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 15 5-chloro-*N*-(2-oxo-1-{2-oxo-2-(pyridin-4-ylamino)ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(1-methyl-1*H*-pyrazol-5-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(1,3-dimethyl-1*H*-pyrazol-5-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 20 5-chloro-*N*-(2-oxo-1-{2-oxo-2-[(pyrazin-2-ylmethyl)amino]ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(6-fluoropyridin-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 25 5-chloro-*N*-(1-{2-[(2-hydroxypyrimidin-4-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(2-oxo-1-{2-oxo-2-(pyrimidin-4-ylamino)ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(1-ethyl-1*H*-pyrazol-5-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 30 5-chloro-*N*-(2-oxo-1-{2-oxo-2-[(5-oxo-4,5-dihydro-1*H*-pyrazol-3-yl)amino]ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;

-83-

- 5-chloro-*N*-(1-{2-[(4-hydroxypyrimidin-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(3-methylpyridin-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5 5-chloro-*N*-(1-{2-[(6-chloropyridazin-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(1*H*-imidazol-2-ylmethyl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(1-methyl-1*H*-pyrazol-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 10 5-chloro-*N*-(1-{2-[(3-ethyl-1*H*-pyrazol-5-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(3-ethyl-1*H*-pyrazol-5-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 15 5-chloro-*N*-(1-{2-[(5-fluoropyridin-2-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- N*-(1-{2-[(6-bromopyridin-3-yl)amino]-2-oxoethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-5-chloro-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(2-hydroxyethyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;
- 20 5-chloro-*N*-(1-{2-[(2,2-dimethyl-1,3-dioxan-5-yl)methyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(3-hydroxy-2-(hydroxymethyl)propyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 25 5-chloro-*N*-(1-{2-[(2,3-dihydroxypropyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(3-hydroxy-2-oxopropyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(1-{2-[(2*R*)-2,3-dihydroxypropyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 30 5-chloro-*N*-(1-{2-[(methylsulfonyl)amino]ethyl}-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;

-84-

- N*-{1-[2-(acetylamino)ethyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl}-5-chloro-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(2-oxo-1-{2-[(trifluoroacetyl)amino]ethyl}-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-[1-(3-hydroxypropyl)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide;
- N*-{1-[(2*Z*)-2-amino-2-(hydroxyimino)ethyl]-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl}-5-chloro-1*H*-indole-2-carboxamide;
- 5-chloro-*N*-(6-fluoro-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl)-1*H*-indole-2-carboxamide; and
- 10 5-chloro-*N*-[6-(methyloxy)-2-oxo-1,2,3,4-tetrahydroquinolin-3-yl]-1*H*-indole-2-carboxamide; or a pharmaceutically acceptable salt or an in-vivo hydrolysable ester thereof.

10. A pharmaceutical composition which comprises a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of claims 1 to 9 in association with a pharmaceutically-acceptable diluent or carrier.

11. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of claims 1 to 9, for use in a method of treatment of a warm-blooded animal such as man by therapy.

20

12. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of claims 1 to 9, for use as a medicament.


13. A compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of claims 1 to 9, for use as a medicament in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

14. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or in-vivo hydrolysable ester thereof, as claimed in any one of claims 1 to 9, in the manufacture of a medicament for use in the treatment of type 2 diabetes, insulin resistance, syndrome X, hyperinsulinaemia, hyperglucagonaemia, cardiac ischaemia or obesity in a warm-blooded animal such as man.

15. The use of a compound of the formula (1), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as claimed in any one of claims 1 to 9, in the manufacture of a medicament for use in the treatment of type 2 diabetes in a warm-blooded animal such as
 5 man.

16. A process for the preparation of a compound of formula (1) as claimed in claim 1, which process comprises:


reacting an acid of the formula (2):



10

(2)

or an activated derivative thereof; with an amine of formula (3):



(3)

15 and thereafter if necessary:

- i) converting a compound of the formula (1) into another compound of the formula (1);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

20

25